

Partial Derivatives of Observables with Respect to Two-Body Orbital Elements

Marc A. Murison
U.S. Naval Observatory, Washington, DC
murison@usno.navy.mil

18 December, 2006

Abstract

We calculate the partial derivatives of the observable quantities $\{\alpha, \delta, r, \dot{r}\}$, where α is the right ascension and δ is the declination, with respect to the two-body orbital elements $\{a, e, \iota, \Omega, \omega, E\}$, where E is the eccentric anomaly. These observables are the only ones available, at least in principle, for the determination of orbital elements of orbiting bodies. We avoid the complexity of expressing the observables directly in terms of orbital elements by instead expressing them in Cartesian coordinates and then invoking the chain rule for derivatives.

Subject headings: celestial mechanics—two-body problem—astrometry—observables

The XML version of this document is available on the web at
http://www.alpheratz.net/murison/dynamics/twobody/twobody_partials.xml

The PDF version of this document is available on the web at
http://www.alpheratz.net/murison/dynamics/twobody/twobody_partials.pdf

1	Components of the Two-Body State Vector	3
1.1	The Position Vector	3
1.2	Scalar Distance	4
1.3	Radial Velocity	5
1.4	Velocity Vector	5
2	Partial Derivatives of the State Vector	6
2.1	Position	6
2.2	Velocity	7
3	Observables and the State Vector	8
3.1	Astrometric Angles	8
3.2	Scalar Distance	9
3.3	Radial Velocity	9
4	Partials of the Observables	9
4.1	Astrometric Angles	10
4.2	Scalar Distance	10
4.3	Radial Velocity	10
4.4	Evaluation of the Partial Derivatives	10
5	Appendix: Explicit State Vector Derivatives	11

1 Components of the Two-Body State Vector

1.1 The Position Vector




Figure 1: The orbital elements and angular momentum vector.

Figure 1 shows an orbit with the standard set of orbital elements $\{a, e, \iota, \omega, \Omega, \theta\}$, where θ is the true anomaly and \vec{h} is the angular momentum (per mass). To reduce expression complexity, we shall use the eccentric anomaly E as the fast angular variable instead of the true anomaly. See Figure 2.

An unperturbed two-body elliptical orbit radius vector can be written

$$\vec{r}(E) = a\mathcal{Q}(\Omega, \iota, \omega) \cdot \vec{q}(E) \quad (1)$$

where a is the semimajor axis, e the orbital eccentricity, ι the orbital inclination, Ω the longitude of the ascending node, and ω the argument of pericenter. The position/shape vector is

$$\vec{q}(E) = \langle \cos E - e, \sqrt{1 - e^2} \sin E, 0 \rangle \quad (2)$$

and the orbit orientation matrix is

$$\mathcal{Q}(\Omega, \iota, \omega) = \begin{bmatrix} \cos\Omega\cos\omega - \sin\Omega\sin\omega\cos\iota & -\cos\Omega\sin\omega - \sin\Omega\cos\omega\cos\iota & \sin\Omega\sin\iota \\ \sin\Omega\cos\omega + \cos\Omega\sin\omega\cos\iota & -\sin\Omega\sin\omega + \cos\Omega\cos\omega\cos\iota & -\cos\Omega\sin\iota \\ \sin\omega\sin\iota & \cos\omega\sin\iota & \cos\iota \end{bmatrix} \quad (3)$$




Figure 2: The geometric relation between true anomaly θ and eccentric anomaly E .

where

$$\mathcal{Q}(\Omega, \iota, \omega) = \mathcal{R}_z(\Omega) \cdot \mathcal{R}_x(\iota) \cdot \mathcal{R}_z(\omega) \quad (4)$$

is the composition of the rotation matrices

$$\mathcal{R}_x(\psi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos\psi & -\sin\psi \\ 0 & \sin\psi & \cos\psi \end{bmatrix} \quad (5)$$

$$\mathcal{R}_z(\psi) = \begin{bmatrix} \cos\psi & -\sin\psi & 0 \\ \sin\psi & \cos\psi & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (6)$$

The matrices (5) and (6) individually rotate a vector counterclockwise around the respective x or z coordinate axis of an inertial frame. (Alternatively, one may interpret these as rotating the coordinate frame clockwise about the respective axes.)

The formulation (1) cleanly separates the orbit scale a , the shape of the orbit plus the position of the particle at a given time $\vec{q}(t)$ (with $E(t)$ obtained via the Kepler equation), and the orientation of the orbit in space $\mathcal{Q}(\Omega, \iota, \omega)$.

1.2 Scalar Distance

The matrix $\mathcal{Q}(\Omega, \iota, \omega)$ is orthogonal, since it is composed of orthogonal matrices. Hence, $\mathcal{Q}(\Omega, \iota, \omega)^\top \cdot \mathcal{Q}(\Omega, \iota, \omega) = \mathcal{I}$, where \mathcal{I} is the identity matrix. The magnitude of the radius

vector can therefore formally be derived from (1) as

$$\begin{aligned}
r^2 &\equiv \vec{r}^\top \cdot \vec{r} \\
&= a^2 (\vec{q}^\top \cdot \mathcal{Q}^\top) \cdot (\mathcal{Q} \cdot \vec{q}) \\
&= a^2 \vec{q}^\top \cdot (\mathcal{Q}^\top \cdot \mathcal{Q}) \cdot \vec{q} \\
&= a^2 (\vec{q}^\top \cdot \mathcal{I} \cdot \vec{q}) \\
&= a^2 (\vec{q}^\top \cdot \vec{q}) \\
&= a^2 q^2
\end{aligned} \tag{7}$$

From the definition (2), $q^2 = (1 - e \cos E)^2$. Thus, the scalar distance is

$$r(E) = \sqrt{\vec{r}(E)^\top \cdot \vec{r}(E)} = a \sqrt{\vec{q}(E)^\top \cdot \vec{q}(E)} = a(1 - e \cos E) \tag{8}$$

1.3 Radial Velocity

The radial velocity is the time derivative of the magnitude of the position vector. If we write the velocity vector in radial and perpendicular parts,

$$\vec{v} = \dot{r} \hat{r} + r \dot{\theta} \hat{\theta} \tag{9}$$

One immediately sees that the projection onto the line of sight of an observer at the coordinate origin is

$$\vec{v} \cdot \hat{r} = \dot{r} \tag{10}$$

We shall use (8) to calculate \dot{r} . We first need \dot{E} . Thus, take the time derivative of the Kepler equation

$$E - e \sin E = M = n(t - \tau) \tag{11}$$

where n is the mean motion and τ is the time of pericenter passage, to get

$$\dot{E}(1 - e \cos E) = n \tag{12}$$

which by (8) is just

$$\dot{E} = \frac{na}{r} \tag{13}$$

Differentiating (8), we find our result,

$$\dot{r} = \frac{na^2}{r} e \sin E = \frac{na}{1 - e \cos E} e \sin E \tag{14}$$

1.4 Velocity Vector

The two-body velocity is, by differentiating (1),

$$\vec{v} = a \mathcal{Q}(\Omega, \iota, \omega) \cdot \frac{\partial \vec{q}(E)}{\partial E} \dot{E} \tag{15}$$

From (2), (8), and (13) follows the result

$$\begin{aligned}\vec{v} &= \frac{na}{1 - e\cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \frac{\partial \vec{q}(E)}{\partial E} \\ &= \frac{na}{1 - e\cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \begin{bmatrix} -\sin E \\ \sqrt{1 - e^2} \cos E \\ 0 \end{bmatrix}\end{aligned}\quad (16)$$

2 Partial Derivatives of the State Vector

2.1 Position

From (1) it is straightforward to calculate the partial derivatives of the position vector with respect to the orbital parameters $\{a, e, \iota, \Omega, \omega, E\}$. First, though, recall that $\mu = n^2 a^3$ (Kepler's third law), where $\mu = G(m_1 + m_2)$ conveniently hides most of the physical units. Thus, for the derivative with respect to semimajor axis, we see by the Kepler equation, (11), that the eccentric anomaly is an explicit function of semimajor axis, so that

$$\frac{\partial E}{\partial a} = -\frac{3n}{2r}(t - \tau) \quad (17)$$

We then find

$$\begin{aligned}\frac{\partial \vec{r}}{\partial E} &= a \mathcal{Q}(\Omega, \iota, \omega) \cdot \frac{\partial \vec{q}(E)}{\partial E} \\ &= a \mathcal{Q}(\Omega, \iota, \omega) \cdot \langle -\sin E, \sqrt{1 - e^2} \cos E, 0 \rangle\end{aligned}\quad (18)$$

$$\begin{aligned}\frac{\partial \vec{r}}{\partial a} &= \mathcal{Q}(\Omega, \iota, \omega) \cdot \left(\vec{q}(E) + a \frac{\partial \vec{q}(E)}{\partial E} \frac{\partial E}{\partial a} \right) \\ &= \frac{\vec{r}}{a} - \frac{3na}{2r}(t - \tau) \mathcal{Q}(\Omega, \iota, \omega) \cdot \langle -\sin E, \sqrt{1 - e^2} \cos E, 0 \rangle\end{aligned}\quad (19)$$

$$\begin{aligned}\frac{\partial \vec{r}}{\partial e} &= a \mathcal{Q}(\Omega, \iota, \omega) \cdot \frac{\partial \vec{q}(E)}{\partial e} \\ &= -a \mathcal{Q}(\Omega, \iota, \omega) \cdot \left\langle 1, \frac{e \sin E}{\sqrt{1 - e^2}}, 0 \right\rangle\end{aligned}\quad (20)$$

$$\frac{\partial \vec{r}}{\partial \{\Omega, \iota, \omega\}} = a \frac{\partial \mathcal{Q}(\Omega, \iota, \omega)}{\partial \{\Omega, \iota, \omega\}} \cdot \vec{q}(E) \quad (21)$$

2.2 Velocity

For the quantity na that appears in the velocity terms (cf. eq. (16)), we have

$$na = \sqrt{\frac{\mu}{a}} \quad (22)$$

One must be careful to not naively take $\partial(na)/\partial a = n$. Right away, something must be wrong with that, since na is a velocity (the circular velocity of an orbit with radius a), and physical intuition indicates that an increase in semimajor axis should cause na to decrease. Instead we must take the derivative of the right-hand side of (22). Hence,

$$\frac{\partial(na)}{\partial a} = \frac{\partial}{\partial a} \sqrt{\frac{\mu}{a}} = -\frac{1}{2} \sqrt{\frac{\mu}{a^3}} = -\frac{1}{2}n \quad (23)$$

One must also remember the dependence of E on semimajor axis in the derivative of \vec{v} with respect to a (cf. eq. (17)). The partials of the velocity vector (16) then become

$$\begin{aligned} \frac{\partial \vec{v}}{\partial a} &= -\frac{1}{2} \frac{n}{1 - e \cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \frac{\partial \vec{q}(E)}{\partial E} + \frac{na}{1 - e \cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \frac{\partial^2 \vec{q}(E)}{\partial E^2} \frac{\partial E}{\partial a} \\ &= -\frac{1}{2} \frac{\vec{v}}{a} + \frac{3}{2} (t - \tau) \left(\frac{na}{r} \right)^2 \mathcal{Q}(\Omega, \iota, \omega) \cdot \begin{bmatrix} \cos E \\ \sqrt{1 - e^2} \sin E \\ 0 \end{bmatrix} \end{aligned} \quad (24)$$

$$\begin{aligned} \frac{\partial \vec{v}}{\partial e} &= \frac{na}{1 - e \cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \left(\frac{\cos E}{1 - e \cos E} \frac{\partial \vec{q}(E)}{\partial E} + \frac{\partial^2 \vec{q}(E)}{\partial E \partial e} \right) \\ &= \frac{na}{1 - e \cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \left(\frac{\cos E}{1 - e \cos E} \begin{bmatrix} -\sin E \\ \sqrt{1 - e^2} \cos E \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ \frac{-e \cos E}{\sqrt{1 - e^2}} \\ 0 \end{bmatrix} \right) \end{aligned} \quad (25)$$

$$\begin{aligned} \frac{\partial \vec{v}}{\partial \{\Omega, \iota, \omega\}} &= \frac{na}{1 - e \cos E} \frac{\partial \mathcal{Q}(\Omega, \iota, \omega)}{\partial \{\Omega, \iota, \omega\}} \cdot \frac{\partial \vec{q}(E)}{\partial E} \\ &= \frac{na}{1 - e \cos E} \frac{\partial \mathcal{Q}(\Omega, \iota, \omega)}{\partial \{\Omega, \iota, \omega\}} \cdot \langle -\sin E, \sqrt{1 - e^2} \cos E, 0 \rangle \end{aligned} \quad (26)$$

$$\begin{aligned} \frac{\partial \vec{v}}{\partial E} &= \frac{-na}{1 - e \cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \left(\frac{e \sin E}{1 - e \cos E} \frac{\partial \vec{q}(E)}{\partial E} - \frac{\partial^2 \vec{q}(E)}{\partial E^2} \right) \\ &= \frac{-na}{1 - e \cos E} \mathcal{Q}(\Omega, \iota, \omega) \cdot \left(\frac{e \sin E}{1 - e \cos E} \begin{bmatrix} -\sin E \\ \sqrt{1 - e^2} \cos E \\ 0 \end{bmatrix} + \begin{bmatrix} \cos E \\ \sqrt{1 - e^2} \sin E \\ 0 \end{bmatrix} \right) \end{aligned} \quad (27)$$

For completeness, in the Appendix we explicitly expand eqs. (18)-(21) and (24)-(27) to component form. A file containing C language subroutines that numerically calculate the derivatives in eqs. (18)-(21) and (24)-(27), optimized with respect to cpu time, is available at http://www.alpheratz.net/murison/dynamics/twobody/twobody_partials.c.

3 Observables and the State Vector

When observing orbiting bodies in the Solar System, we potentially have up to four observable quantities. The angular position on the plane of the sky is specified by two observables, say right ascension and declination, and is provided by classical astrometry. The radial velocity can be had from spectroscopy or doppler radar. Finally, the scalar distance can be found by radar ranging. In Section 4, we will calculate the partial derivatives of these observables with respect to the orbital elements. First, though, it will be convenient to determine the partial derivatives of the observables with respect to the Cartesian state vector

$$\vec{S} = \langle x, y, z, \dot{x}, \dot{y}, \dot{z} \rangle \tag{28}$$

as an intermediate step, and then invoke the chain rule for derivatives. This approach is more useful because it would be difficult to express the observables directly as functions of the orbital elements. Thus, we need to relate explicitly the observables to the components of \vec{S} .

3.1 Astrometric Angles




Figure 3: Right ascension α and declination δ .

From Figure 3 we see that

$$\begin{aligned}x &= r \cos \delta \cos \alpha \\y &= r \cos \delta \sin \alpha \\z &= r \sin \delta\end{aligned}\tag{29}$$

where α is the right ascension and δ is the declination. We can write (29) compactly as the unit vector

$$\frac{\vec{r}}{r} = \vec{F}(\alpha, \delta)\tag{30}$$

The inverse is then

$$\langle \alpha, \delta \rangle = \vec{G}(\vec{r})\tag{31}$$

From (29) (or by inspection of Figure 3), eq. (31) becomes

$$\begin{aligned}\sin \delta &= \frac{z}{r} \\ \sin \alpha &= \frac{y/r}{\sqrt{1 - (z/r)^2}} = \frac{y}{\sqrt{x^2 + y^2}} \\ \cos \alpha &= \frac{x/r}{\sqrt{1 - (z/r)^2}} = \frac{x}{\sqrt{x^2 + y^2}}\end{aligned}\tag{32}$$

We therefore can easily convert back and forth between the position vector and the angular observables, and from (32) we can calculate the Cartesian derivatives of $\{\alpha, \delta\}$.

3.2 Scalar Distance

The scalar distance is just

$$r = \sqrt{x^2 + y^2 + z^2}\tag{33}$$

3.3 Radial Velocity

The radial velocity can be written

$$\dot{r} = \vec{v} \cdot \hat{r} = \frac{\vec{v} \cdot \vec{r}}{r} = \frac{x\dot{x} + y\dot{y} + z\dot{z}}{\sqrt{x^2 + y^2 + z^2}}\tag{34}$$

4 Partial derivatives of the Observables

We shall now calculate the partials of the observables with respect to the Cartesian state vector components, $\frac{\partial \varphi}{\partial s_k}$. Once that is done, then eq. (40) in section 4.4 can be evaluated for the partial derivatives of each observable with respect to each of the orbital elements.

4.1 Astrometric Angles

From eqs. (32), we find

$$\begin{aligned}\frac{\partial\alpha}{\partial x} &= -\frac{y}{x^2+y^2} \\ \frac{\partial\alpha}{\partial y} &= \frac{x}{x^2+y^2} \\ \frac{\partial\alpha}{\partial z} &= 0\end{aligned}\tag{35}$$

and

$$\begin{aligned}\frac{\partial\delta}{\partial x} &= -\frac{xz}{r^2}\frac{1}{\sqrt{x^2+y^2}} \\ \frac{\partial\delta}{\partial y} &= -\frac{yz}{r^2}\frac{1}{\sqrt{x^2+y^2}} \\ \frac{\partial\delta}{\partial z} &= \frac{\sqrt{x^2+y^2}}{r^2}\end{aligned}\tag{36}$$

4.2 Scalar Distance

From (33) we have

$$\frac{\partial r}{\partial\{x,y,z\}} = \frac{\{x,y,z\}}{r}\tag{37}$$

4.3 Radial Velocity

From (34) we calculate

$$\frac{\partial\dot{r}}{\partial\{x,y,z\}} = \frac{\{\dot{x},\dot{y},\dot{z}\}}{r} - \frac{\dot{r}}{r^2}\{x,y,z\}\tag{38}$$

and

$$\frac{\partial\dot{r}}{\partial\{\dot{x},\dot{y},\dot{z}\}} = \frac{\{x,y,z\}}{r}\tag{39}$$

4.4 Evaluation of the Partial Derivatives

Let $p \in \{a, e, \iota, \Omega, \omega, E\}$ be an orbital parameter, and let $\varphi \in \{\alpha, \delta, r, \dot{r}\}$ be an observable. Then the partial derivatives of the observables with respect to the orbital elements can be written

$$\frac{\partial\varphi}{\partial p} = \frac{\partial\varphi}{\partial\vec{S}} \cdot \frac{\partial\vec{S}}{\partial p} = \sum_{k=1}^6 \frac{\partial\varphi}{\partial s_k} \frac{\partial s_k}{\partial p}\tag{40}$$

where $s_k = \left[\vec{S} \right]_k$ is the k^{th} component of the Cartesian state vector, eq. (28). Equation (40) is the fundamental relation. The derivatives $\frac{\partial s_k}{\partial p}$ are given by eqs. (18)-(21) and (24)-(27), and the $\frac{\partial \varphi}{\partial s_k}$ are given by eqs. (35)-(40).

5 Appendix: Explicit State Vector Derivatives

Recall that the state vector in Cartesian components is

$$\vec{S} = \langle x, y, z, \dot{x}, \dot{y}, \dot{z} \rangle \quad (41)$$

From eqs. (18)-(21) and (24)-(27) we explicitly calculate the components of the partial derivatives of \vec{S} with respect to the orbital elements $\{a, e, \iota, \Omega, \omega, E\}$. Optimized C code subroutines for calculating these derivatives numerically is available at http://www.alpheratz.net/murison/dynamics/twobody/twobody_partials.c.

$$\begin{aligned} \frac{\partial \vec{S}}{\partial a} = & \begin{bmatrix} -\sqrt{1-e^2}(\cos\Omega\sin\omega + \sin\Omega\cos\omega\cos\iota)\sin E - (\cos E - e)(-\cos\Omega\cos\omega + \sin\Omega\sin\omega\cos\iota) \\ -\sqrt{1-e^2}(\sin\Omega\sin\omega - \cos\Omega\cos\omega\cos\iota)\sin E + (\cos E - e)(\sin\Omega\cos\omega + \cos\Omega\sin\omega\cos\iota) \\ \left[\sqrt{1-e^2}\cos\omega\sin E + (\cos E - e)\sin\omega \right] \sin\iota \\ \frac{1}{2} \frac{n}{1-\epsilon\cos E} \left[\sqrt{1-e^2}(\cos\Omega\sin\omega + \sin\Omega\cos\omega\cos\iota)\cos E - \sin E(-\cos\Omega\cos\omega + \sin\Omega\sin\omega\cos\iota) \right] \\ \frac{1}{2} \frac{n}{1-\epsilon\cos E} \left[-\sqrt{1-e^2}(-\sin\Omega\sin\omega + \cos\Omega\cos\omega\cos\iota)\cos E + \sin E(\sin\Omega\cos\omega + \cos\Omega\sin\omega\cos\iota) \right] \\ \frac{1}{2} \frac{n}{1-\epsilon\cos E} \left(-\sqrt{1-e^2}\cos\omega\cos E + \sin\omega\sin E \right) \sin\iota \end{bmatrix} \\ & + \frac{3}{2} \frac{n(t-\tau)}{1-\epsilon\cos E} \\ & \times \begin{bmatrix} \sqrt{1-e^2}(\cos\Omega\sin\omega + \sin\Omega\cos\omega\cos\iota)\cos E - (-\cos\Omega\cos\omega + \sin\Omega\sin\omega\cos\iota)\sin E \\ \sqrt{1-e^2}(\sin\Omega\sin\omega - \cos\Omega\cos\omega\cos\iota)\cos E + (\sin\Omega\cos\omega + \cos\Omega\sin\omega\cos\iota)\sin E \\ \left(-\sqrt{1-e^2}\cos\omega\cos E + \sin\omega\sin E \right) \sin\iota \\ \frac{-n}{(1-\epsilon\cos E)^2} \left[\sqrt{1-e^2}(\cos\Omega\sin\omega + \sin\Omega\cos\omega\cos\iota)\sin E - (\cos E - e)(-\cos\Omega\cos\omega + \sin\Omega\sin\omega\cos\iota) \right] \\ \frac{n}{(1-\epsilon\cos E)^2} \left[\sqrt{1-e^2}(-\sin\Omega\sin\omega + \cos\Omega\cos\omega\cos\iota)\sin E + (\cos E - e)(\sin\Omega\cos\omega + \cos\Omega\sin\omega\cos\iota) \right] \\ \frac{n}{(1-\epsilon\cos E)^2} \left[\sqrt{1-e^2}\cos\omega\sin E + (\cos E - e)\sin\omega \right] \sin\iota \end{bmatrix} \end{aligned} \quad (42)$$

$$\frac{\partial \vec{S}}{\partial e} = \begin{bmatrix} a \left(-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota + \frac{e}{\sqrt{1-e^2}} (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \sin E \right) \\ a \left(-\sin\Omega \cos\omega - \cos\Omega \sin\omega \cos\iota + \frac{e}{\sqrt{1-e^2}} (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \sin E \right) \\ -a \left(\sin\omega + \frac{e}{\sqrt{1-e^2}} \cos\omega \sin E \right) \sin\iota \\ \frac{na \cos E}{(1-e \cos E)^2} \left[(-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \sin E - \frac{\cos E - e}{\sqrt{1-e^2}} (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \right] \\ \frac{na \cos E}{(1-e \cos E)^2} \left[-(\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \sin E - \frac{\cos E - e}{\sqrt{1-e^2}} (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \right] \\ \frac{na \cos E}{(1-e \cos E)^2} \left[-\sin\omega \sin E + \frac{\cos E - e}{\sqrt{1-e^2}} \cos\omega \right] \sin\iota \end{bmatrix} \quad (43)$$

$$\frac{\partial \vec{S}}{\partial \iota} = \begin{bmatrix} a \left[\sqrt{1-e^2} \cos\omega \sin E + (\cos E - e) \sin\omega \right] \sin\Omega \sin\iota \\ -a \left[\sqrt{1-e^2} \cos\omega \sin E + (\cos E - e) \sin\omega \right] \cos\Omega \sin\iota \\ a \left[\sqrt{1-e^2} \cos\omega \sin E + (\cos E - e) \sin\omega \right] \cos\iota \\ \frac{na}{1-e \cos E} \left(\sqrt{1-e^2} \cos\omega \cos E - \sin\omega \sin E \right) \sin\Omega \sin\iota \\ \frac{-na}{1-e \cos E} \left(\sqrt{1-e^2} \cos\omega \cos E - \sin\omega \sin E \right) \cos\Omega \sin\iota \\ \frac{na}{1-e \cos E} \left(\sqrt{1-e^2} \cos\omega \cos E - \sin\omega \sin E \right) \cos\iota \end{bmatrix} \quad (44)$$

$$\frac{\partial \vec{S}}{\partial \Omega} = \begin{bmatrix} a \left[\sqrt{1-e^2} (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \sin E - (\cos E - e) (\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \right] \\ a \left[-\sqrt{1-e^2} (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \sin E - (\cos E - e) (-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \right] \\ 0 \\ \frac{na}{1-e \cos E} \left[\sqrt{1-e^2} (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \cos E + (\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \sin E \right] \\ \frac{na}{1-e \cos E} \left[-\sqrt{1-e^2} (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \cos E + (-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \sin E \right] \\ 0 \end{bmatrix} \quad (45)$$

$$\frac{\partial \vec{S}}{\partial \omega} = \begin{bmatrix} a \left[\sqrt{1-e^2} (-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \sin E - (\cos E - e) (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \right] \\ -a \left[\sqrt{1-e^2} (\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \sin E + (\cos E - e) (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \right] \\ a \left[-\sqrt{1-e^2} \sin\omega \sin E + (\cos E - e) \cos\omega \right] \sin\iota \\ \frac{na}{1-e \cos E} \left[\sqrt{1-e^2} (-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \cos E + (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \sin E \right] \\ \frac{na}{1-e \cos E} \left[-\sqrt{1-e^2} (\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \cos E + (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \sin E \right] \\ \frac{-na}{1-e \cos E} \left(\sqrt{1-e^2} \sin\omega \cos E + \cos\omega \sin E \right) \sin\iota \end{bmatrix} \quad (46)$$

$$\frac{\partial \vec{S}}{\partial E} = \begin{bmatrix} -a \left[\sqrt{1-e^2} (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \cos E - (-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \sin E \right] \\ -a \left[\sqrt{1-e^2} (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \cos E + (\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \sin E \right] \\ a \left(\sqrt{1-e^2} \cos\omega \cos E - \sin\omega \sin E \right) \sin\iota \\ \frac{na}{(1-e\cos E)^2} \left[\sqrt{1-e^2} (\cos\Omega \sin\omega + \sin\Omega \cos\omega \cos\iota) \sin E + (\cos E - e) (-\cos\Omega \cos\omega + \sin\Omega \sin\omega \cos\iota) \right] \\ \frac{na}{(1-e\cos E)^2} \left[\sqrt{1-e^2} (\sin\Omega \sin\omega - \cos\Omega \cos\omega \cos\iota) \sin E - (\cos E - e) (\sin\Omega \cos\omega + \cos\Omega \sin\omega \cos\iota) \right] \\ \frac{-na}{(1-e\cos E)^2} \left[\sqrt{1-e^2} \cos\omega \sin E + (\cos E - e) \sin\omega \right] \sin\iota \end{bmatrix} \quad (47)$$